




LISTING OF CLAIMS


Claims 1 – 30 (canceled)

31. (previously presented) A process for the industrial synthesis of strontium ranelate
5 of formula (I):



and its hydrates,

10 wherein a compound of formula (II) :



wherein R and R', which may be identical or different, each represent linear or branched (C₁-C₆)alkyl,

15 is reacted with strontium hydroxide in an amount greater than or equal to 2 mol per mol of compound of formula (II),

at reflux with water,

for at least 5 hours;

the precipitate obtained is then filtered off while hot;

the cake obtained is washed with boiling water


to yield, after drying of the powder thereby obtained, the compound of formula (I) and its hydrates.

5 **32.** (previously presented) The synthesis process of Claim 31, wherein the amount of water used in the reaction of the compound of formula (II) with strontium hydroxide is greater than or equal to 8 ml per gram of compound of formula (II).

10 **33.** (previously presented) The synthesis process of Claim 31, wherein the amount of strontium hydroxide is from 2 to 2.5 mol per mol of compound of formula (II).


34. (previously presented) The synthesis process of Claim 31, wherein R represents methyl and R' represents methyl or ethyl.

15 **35.** (currently amended) ~~The synthesis process of Claim 31~~ A process for the industrial synthesis of strontium ranelate of formula (I):



20 and its hydrates,

wherein a compound of formula (IV):



wherein R represents linear or branched (C₁-C₆)alkyl,


is reacted with malononitrile of formula (V) :



in methanol,

5 in the presence of morpholine in an amount greater than 0.95 mol per mol of compound of formula (IV),

to yield the compound of formula (VI) :




10 which is then reacted with sulphur in an amount greater than 0.95 mol per mol of compound of formula (IV);


the reaction mixture is then heated at reflux;

and the compound thereby obtained is isolated by precipitation in the presence of water, followed by filtration,

15 to yield the compound of formula (III) :



which is reacted with a compound of formula (VII) :



wherein R' represents linear or branched (C₁-C₆)alkyl,
in the presence of a catalytic amount of a C₈-C₁₀-type quaternary ammonium compound,
and in the presence of potassium carbonate,

at reflux with an organic solvent;


the reaction mixture is subsequently filtered;

the mixture is then concentrated by distillation;

a co-solvent is then added,

and the reaction mixture is cooled and filtered

to yield, after drying of the powder thereby obtained, the compound of formula (II) :



which is reacted with strontium hydroxide according to the process of Claim 31. ~~in an amount greater than or equal to 2 mol per mol of compound of formula (II),~~

~~at reflux with water,~~


~~for at least 5 hours;~~

~~the precipitate obtained is then filtered off while hot;~~

~~the cake obtained is washed with boiling water~~

~~to yield, after drying of the powder thereby obtained, the compound of formula (I) and its hydrates;~~

it being understood that a C₈-C₁₀-type quaternary ammonium compound is a compound of formula (A) or a mixture of compounds of formula (A) :-



wherein ~~R₁ represents (C₁-C₆)alkyl, R₂, R₃ and R₄, which are the same or different, each represent (C₈-C₁₀)alkyl, and X represents a halogen atom.~~

- 5 36. (previously presented) The synthesis process of Claim 35, wherein the amount of methanol used in the synthesis of the compound of formula (III) is from 1 to 3 ml per gram of compound of formula (IV).
37. (previously presented) The synthesis process of Claim 35, wherein the temperature of reaction between the compounds of formulae (IV) and (V) is less than 50°C.
- 10 38. (previously presented) The synthesis process of Claim 35, wherein the refluxing time for the reaction between the compound of formula (VI) and sulphur is between 1 hour 30 minutes and 3 hours.
39. (previously presented) The synthesis process of Claim 35, wherein the amount of potassium carbonate used in the synthesis of the compound of formula (II) is from 2 to
15 3 mol per mol of compound of formula (III).
40. (previously presented) The synthesis process of Claim 35, wherein the amount of compound of formula (VII) is from 2 to 3 mol per mol of compound of formula (III).
41. (previously presented) The synthesis process of Claim 35, wherein the initial volume of organic solvent used in the reaction of the compound of formula (III) with the
20 compound of formula (VII) is from 6 to 12 ml per gram of compound of formula (III).
42. (previously presented) The synthesis process of Claim 35, wherein the organic solvent used in the reaction of the compound of formula (III) with the compound of formula (VII) is acetone or acetonitrile.

43. (previously presented) The synthesis process of Claim 35, wherein the co-solvent used in the isolation of the compound of formula (II) is methanol.

44. (previously presented) The synthesis process of Claim 35, wherein the compound of formula (II) obtained has a chemical purity greater than 98 %.

5 45. (previously presented) The synthesis process of Claim 35, wherein the amount of water used in the reaction of the compound of formula (II) with strontium hydroxide is greater than or equal to 8 ml per gram of compound of formula (II).

46. (previously presented) The synthesis process of Claim 35, wherein the amount of strontium hydroxide is from 2 to 2.5 mol per mol of compound of formula (II).

10 47. (previously presented) The synthesis process of Claim 35, wherein R represents methyl and R' represents methyl or ethyl.